

ΔNewYork Blood Center

IMMUNIZATION OR VACCINATION	DEFER FOR:
Allergy Desensitization	Accept immediately
Anthrax	Accept immediately if feeling well and healthy
Botulinum Toxin ('Botox')	Defer for 24 hours after treatment
BCG	2 weeks
Chicken Pox (Varivax)	4 weeks
Chicken Pox (VZIG) Immune Globulin	12 months
Cholera	Accept immediately if feeling well and healthy
COVID-19	<ul style="list-style-type: none"> • Accept immediately if feeling well and healthy after receiving an mRNA vaccine such as Pfizer or Moderna • 2 weeks if feeling well and healthy after receiving an attenuated vaccine such as AstraZeneca • 2 weeks if feeling well and healthy and vaccine type is unknown or if vaccine is in clinical trials • If the donor was infected with COVID-19 prior to receiving the COVID-19 vaccine, the donor is eligible for CCP for 6 months after their last COVID symptom • Donors who received the vaccine prior to COVID-19 infection are not eligible for CCP donation • Individuals who received an investigational COVID-19 monoclonal antibody therapy as a participant in a clinical trial or received an authorized or licensed COVID-19 monoclonal antibody therapy, are deferred from donating CCP until at least three months after receipt of the therapy. Accept immediately if feeling well and healthy for all other donations.
Diphtheria	Accept immediately if feeling well and healthy
DT Diphtheria \ Tetanus	Accept immediately if feeling well and healthy
DPT Diphtheria \ Pertussis \ Tetanus	Accept immediately if feeling well and healthy
Flu (injection or nasal)	Accept immediately if feeling well and healthy
Gamma Globulin - prophylactic	Accept immediately if feeling well and healthy and given for general prophylaxis (e.g., for travel) and not for exposure
Gamma Globulin - exposure	<ul style="list-style-type: none"> • 120 days (4 months) if given as a precaution to Hepatitis A exposure (i.e., restaurant outbreak). • Evaluate if given for exposure to any other disease.
Hepatitis A - prophylactic	Accept immediately if feeling well and healthy
Hepatitis A – exposure	120 days (4 months)
Hepatitis B - prophylactic	2 weeks
Hepatitis B – exposure	12 months
Hep A + Hep B - prophylactic	2 weeks
Hep A + Hep B - exposure	12 months
HIB, Haemophilus influenzae b	Accept immediately if feeling well and healthy

ΔNewYork Blood Center

IMMUNIZATION OR VACCINATION	DEFER FOR:
Human Papilloma Virus	Accept immediately if feeling well and healthy
IVIG (Intravenous Immune Globulin)	Defer for 12 months.
Japanese Encephalitis	Accept immediately if feeling well and healthy
Lyme	Accept immediately if feeling well and healthy
Measles (Rubeola)	2 weeks
Measles, German (Rubella)	4 weeks
Meningococcus (including Meningitis B)	Accept immediately if feeling well and healthy
MMR (measles, mumps, rubella)	4 weeks
Mumps	2 weeks
Paratyphoid	Accept immediately if feeling well and healthy
Pertussis	Accept immediately if feeling well and healthy
Plague	Accept immediately if feeling well and healthy
Pneumonia	Accept immediately if feeling well and healthy
Polio, oral	2 weeks
Polio, injection	Accept immediately if feeling well and healthy
Rabies	<ul style="list-style-type: none"> • Accept immediately if received for prophylaxis (e.g., animal care worker) and not for exposure • Defer 12 months if due to animal bite
Rocky Mountain Spotted Fever	Accept immediately if feeling well and healthy
Shingles: Zostavax)	4 weeks
Shingles: Shingrix (Zoster Vaccine Recombinant)	Accept immediately if feeling well and healthy
Smallpox without complications	8 weeks
Smallpox with complications	<ul style="list-style-type: none"> • 8 weeks after onset of symptoms or 14 days after symptoms resolve, whichever is longer • Complications include the following: Rash (resembling 'blisters') covering a small or large area of the body; necrosis (tissue death) in the area of vaccination; inflammation of the brain (encephalitis); infection of the cornea (eye); localized or systemic skin reaction in someone with eczema or other chronic skin condition
Tetanus Toxoid (Booster)	Accept immediately if feeling well and healthy
Tetanus and Diphtheria Toxoids	Accept immediately if feeling well and healthy
Typhoid Injection	Accept immediately if feeling well and healthy
Typhoid Oral	2 weeks
Typhus, Paratyphoid	Accept immediately if feeling well and healthy
Yellow Fever	2 weeks